



Michael S. Day, MD, MPhil

WellSpan Orthopedics

120 North Seventh Street, Chambersburg, PA 17201, Suite 101

Tel: (717) 263-1220; Fax: (717) 263-6255

POST-OPERATIVE INSTRUCTIONS

Shoulder Arthroscopy, Lysis of Adhesions, Manipulation Under Anesthesia

Activity

- Try and rest the first few days following surgery. Wear your **slings** for comfort for the first 2-3 days following surgery. You will be directed by Dr. Day as to when you may discontinue your sling, but generally most patients are able to wean out of the sling by the first week.
- With the sling, you may be more comfortable sleeping in a seated position (i.e. recliner) with a pillow under your forearm for support.
- **Ice** or the **cryo-cuff** can be used as much as possible for the first 3-4 weeks to help decrease swelling.
- You may remove your arm from the sling 4-5 times a day and begin the enclosed **Pendulum & Codman** exercises. Using gravity and momentum gently move your arm in small, slow circles for approximately 5-10 minutes.
- You may also gently bend and extend your elbow to prevent stiffness.
- Do not lift any objects greater than 1-2 pounds for the first 4-6 weeks
- Driving is not recommended for the 1st 2 weeks following surgery and contraindicated when taking narcotic pain medication.
- Return to (sedentary) work or school the day after surgery if pain is tolerable. Return to heavy labor will be determined by Dr. Day.

Dressings

- Remove the outer dressing 48 hours after surgery. Clean the surrounding skin with sterile water or rubbing alcohol and cover the sutures loosely with band-aids or a clean dressing. Be sure not to vigorously rub over any of the incision sites as they are still healing.
- Small amounts of bloody drainage, numbness at incision site, swelling, and bruising are normal findings following surgery.
- Do not use any ointments under band-aids.

Showering/Bathing:

- You may shower after 48 hours after returning home. You can allow the shower water to run down the incision. There is no need to cover the wound while in the shower.
- You should not scrub the incision. Be sure to pat the incision dry with a towel after showering.
- If wound drainage is noted, you should **sponge bathe instead of showering** until the drainage resolves.
- No submersion in water, including a bath, Jacuzzi, lake or swimming pool for six weeks and until given clearance by Dr. Day.
- You may remove the sling to shower, but be careful to avoid any sudden movements or heavy lifting

Medication

- You have been prescribed one of the following **pain medications**:
 - Oxycodone– Please take 1-2 tabs every 4-6 hours as needed for severe pain.
- Post-operative pain tends to be the most pronounced the first 2-3 days following surgery. Please take Tylenol, 1000mg every 6 hours for the first 3 days as this has significantly less side-effects compared to narcotic pain medication. **Do not exceed 4,000 mg acetaminophen in 24 hours from all sources.**
- You may have also been given a prescription for Toradol- take 1 tablet every 6 hours for the first 2 days.
- You have also been given a prescription for Gabapentin (Neurontin) to take one tablet at bedtime as needed for pain.
- You may have also been given a prescription for Aspirin to take 1 tablet (81mg) daily with breakfast for 30 days for DVT/blood clot prophylaxis.
- You may also have been given a prescription for Zofran, an anti-nausea medication. Please take this medication as needed for nausea
- If you have any side effects (i.e. Nausea, rash, trouble breathing) from medication discontinue its use and call our office.
- Excessive use of pain medication can lead to an increased incidence of side effects and long- term dependency issues. Please take the pain medication as directed and understand that the medications are meant to minimize pain in the initial post-operative phase but may not relieve pain completely.



Physical Therapy

- Schedule Physical Therapy to begin approximately the day after surgery. A physical therapy prescription is enclosed and may be given to your therapist at your first visit.
- A list of home exercises has been enclosed for you to do during your first week following surgery.

Follow Up

- You have likely already been scheduled for your first post-operative visit. If you have not received an appointment, please contact the office to schedule an appointment to be seen approximately 14 days after your surgery.
- You will be seen by Dr. Day or his Physician Assistant for your first post-operative visit to review your intra-operative findings and to go over any questions you may have.
- Sutures will also be removed at that time.
- Call the office immediately if you develop a fever (>101.5), chills, chest pain, shortness of breath, excessive incisional drainage, or persistent upper extremity numbness.
- If you have any questions or concerns: please feel free to call the office.

Home Exercises



Pendulums

Stand supporting yourself on a table with your good arm. Let your other arm hang down completely relaxed and, **by moving your trunk only**, slowly allow your arm to move in a circular motion both clockwise and counter clockwise. Keep your circles within the width of your feet.

Codman's

Stand supporting yourself on a table with your good arm. Let your other arm hang down completely relaxed and slowly move it forward & back as well as side to side motion.

*** All exercises to be done for 5-10 minutes, 4-5 times a day.**

Home Exercises (cont'd)

Elbow Curls:

Come out of your sling. Without any extra weight, use your good hand to help bring your other hand towards your shoulder by bending your elbow. Then slowly lower your arm back to a straight position. Then repeat.

Wrist Curls:

While in and out of your sling, make a light fist and move your operative wrist in an up and down movement slowly like you are knocking on a door.

Ball Squeezes:

While in and out of your sling, squeeze the foam ball to exercise the hand, fingers, and wrist muscles. This exercise can be very effective to help promote good circulation and prevent excessive swelling.

*** All exercises to be done for 5-10 minutes, 4-5 times a day.**



The Peak of Good Health

Michael S. Day, MD, MPhil

Summit Orthopedic Group

120 North Seventh Street, Chambersburg, PA 17201, Suite 101

Tel: (717) 263-1220; Fax: (717) 263-6255

https://www.summithealth.org/services/orthopedics

Patient Name: _____

Date: _____

Treatment Frequency 5 x week

Duration of Therapy Prescription: 2 weeks

Treatment Frequency 3 x week

Duration of Therapy Prescription: 4 weeks

PHYSICAL THERAPY PRESCRIPTION:

SHOULDER ARTHROSCOPY WITH CAPSULAR RELEASE AND MANIPULATION

WEEKS 1 - 6

- Anti-Inflammatory Modalities (Ice!) – as needed
- Range of motion exercises all planes to tolerance (ABD, FF, ADD, ER, IR)!
- Codman's, pendulums, pulleys, cane assistive ROM
- INSTRUCT HOME PROGRAM – TO BE DONE DAILY.
- Biceps / triceps isotonic /Elbow & Wrist motion exercises
- Scapular stabilization exercises
- Cardiovascular training as tolerated (Running, bike)
- Anterior and Posterior capsular stretch after warm-up
- Recommend 4-5 visits per week during the first two weeks postoperatively; then 3x per week
- Shoulder CPM: (If box checked will be using a home CPM in addition to PT to restore shoulder motion)
- Special: _____

WEEKS 6- 12

- Range of motion exercises all planes to tolerance (ABD, FF, ADD, ER, IR)!
- Continue upper extremity PRE's
- Continue scapular stabilization / strengthening exercises
- IR / ER isotonic exercises below horizontal (emphasize eccentrics)
- Biceps PRE's
- Continue with shoulder and neck flexibility exercises
- Modalities PRN
- Functional activities begin week 6 (ADL's, Sports)

Please Send Progress notes

Physician's Signature: _____

(I have medically prescribed the above treatments)

Michael S. Day, MD