



Michael S. Day, MD, MPhil

WellSpan Orthopedics

120 North Seventh Street, Chambersburg, PA 17201, Suite 101

Tel: (717) 263-1220; Fax: (717) 263-6255

POST-OPERATIVE INSTRUCTIONS

Ulnar Nerve Decompression

Activity

- Rest the first few days following surgery. When sitting or lying down be sure to keep your arm elevated above the level of your heart to help decrease swelling.
- **Ice** can be used as much as possible for the first 3-4 weeks to help decrease swelling. Ice should be used frequently for 20 minutes at a time.
- Wiggle your exposed fingers regularly to help with circulation. You may use your opposite hand to assist in these exercises.
- Please avoid high force/resistance grasping using your operative hand. Avoid lifting objects heavier than one pound for one month following surgery.
- Driving is not recommended when taking narcotic pain medication.
- Return to (sedentary) work or school the day after surgery if pain is tolerable. Return to heavy labor or excessive ambulatory activities will be determined by Dr. Montgomery.

Dressings

- A large bulky splint has been placed on your elbow/wrist and is designed to feel “snug”.
- This splint is to be left in place for _____.
- The post-operative sling that you were given at the time of surgery is to be used for comfort and does not have to be worn at all times.
- Keep your dressing clean and dry until your first follow-up visit. If you wish to shower, plastic bags, saran wrap, and waterproof dressings can all be used to keep the area dry. **DO NOT SOAK** your knee in a bath tub.

Medication

- You have been prescribed one of the following **pain medications**:
 - Oxycodone – Please take 1 tab every 4-6 hours as needed for pain.
 - Ultram – Please take 1 tab every 6 hours as needed for breakthrough pain
- As the pain gradually improves you may transition over to Extra-Strength Tylenol which is a strong non-narcotic alternative for pain.

- If you have any side effects (i.e. Nausea, rash, trouble breathing) from medication discontinue its use and call our office.

Physical Therapy

- The need for a timing of outpatient physical therapy will be determined by Dr. Montgomery and discussed at your first post-operative visit.
- Please perform your prescribed home exercises to help facilitate early shoulder range of motion and to prevent stiffness in your elbow and wrist.

Follow Up

- You have likely already been scheduled for your first post-operative visit. If you have not received an appointment please contact the office to schedule an appointment 10-14 days after your surgery.
- You will be seen by Dr. Montgomery or his Physician Assistant for your first post-operative visit to review your intra-operative findings and to go over any questions you may have.
- Sutures will also be removed at that time.
- Call the office immediately if you develop a fever (>101.5), chills, excessive incision drainage, calf pain, or persistent upper extremity numbness.
- If you have any questions or concerns please feel free to call the office.